

## Total cordial labeling of corona product of paths and second power of fan graph

Shokry Nada <sup>a</sup>, Atef Abd El-hay <sup>b</sup>, Ashraf Elrokh <sup>c</sup>

<sup>a,b,c</sup>Dept. of Math., Faculty of Science, Menoufia University, Shebeen Elkom, Egypt.

E mail address: <sup>a</sup>shokrynada@yahoo.com, <sup>b</sup>atef\_1992@yahoo.com and <sup>c</sup>ashraf.hefnawy68@yahoo.com

**Abstract:** A graph is called total cordial if it has a 0 – 1 labeling such that the total number of vertices and edges labelled with ones and zeros differ by at most one. In this paper, we contribute some new results on Total Product cordial labeling and investigate necessary and sufficient conditions of the corona product between paths and second power of fan graphs to be total cordial.

**Keywords:** Corona operation, Second power, Total cordial, Path graph, Fan graph.

### 1. Introduction

Graph labeling is one of graph theory's oldest issues. It is the assignment of numbers to vertices, edges, or both under specific condition. Many authors have been interested in graph labeling [1, 3, 4]. Gallian [2] has published an effective survey on whole graph labeling and its applications. The notion of product cordial labeling was introduced in 2004 [6], and it's proved that trees, unicyclic graphs of odd order, triangular snakes, helms, and unions of two path graphs are product cordial.

M. Sundaram, R. Ponraj, and S. Somasundaram introduced a new type of graph labeling known as total product cordial labeling and investigated the total product cordial behavior of some standard graphs [7]. Suppose that  $G = (V, E)$  is a graph, where  $V$  is the set of its vertices and  $E$  is the set of its edges. A mapping  $f: V \rightarrow \{0,1\}$  is called binary vertex labeling of  $G$  and  $f(u)$  denotes the label of vertex  $u$  of  $G$  under  $f$ . A binary vertex labeling of a graph  $G$  with the induced edge labeling  $f^*: E \rightarrow \{0,1\}$  defined by  $f^*(uv) = (f(u) + f(v)) \bmod 2$  is called Total cordial labeling if  $|(v_0 + e_0) - (v_1 + e_1)| \leq 1$ , where  $v_0$  and  $v_1$  are the numbers of vertices labeled by 0 and 1, respectively, and  $e_0$  and  $e_1$  are the corresponding numbers of edges. A graph with a Total cordial labeling defined on it is called Total cordial.

**Definition 1.** A second power of a fan  $F_n^2$  is the graph obtained from the join of the second power of a path  $P_n^2$  and a null graph  $N_1$ , i.e.  $F_n^2 = P_n^2 + N_1$ . So the order of  $F_n^2$  is  $n + 1$  and its size is  $3n - 3$ , in particular  $F_1^2 \equiv P_2, F_2^2 \equiv C_3$  and  $F_3^2 \equiv K_4$ .

**Definition 2.** The corona  $G \odot H$  of two graphs  $G$  (with  $n_1$  vertices and  $m_1$  edges) and  $H$  (with  $n_2$  vertices and  $m_2$  edges) is the graph denoted by  $G \odot H$  and is obtained by taking one copy of  $G$  and  $n_1$  copies of  $H$ , and then joining the  $i^{th}$  vertex of  $G$  with an edge to every vertex in the  $i^{th}$  copy of  $H$  [17]. It follows from the definition of the corona that  $G \odot H$  has  $n_1 + n_1 \cdot n_2$  vertices and  $m_1 + n_1 \cdot m_2 + n_1 \cdot n_2$  edges.

It is easy to see that  $G \odot H$  is not in general isomorphic to  $H \odot G$ .

In this paper we proposed the corona  $P_m \odot F_n^2$  and show that is Total cordial for all  $m \geq 1$  and  $n \geq 4$ .

### 2. Terminologies and Notations

A path with  $m$  vertices and  $m - 1$  edges is denoted by  $P_m$ , and second power of fan graph has  $n + 1$  vertices and  $3n - 3$  edges is denoted by  $F_n^2$ . We let  $M_r$  denote the labeling 0101 ... 01, zero-one repeated  $r$ -times if  $r$  is even and 0101 ... 010 if  $r$  is odd; for example,  $M_6 = 010101$  and  $M_5 = 01010$ . We let  $M'_{2r}$  denote the labeling 1010 ... 10. Sometimes, we modify the labeling  $M_r$  or  $M'_r$  by adding symbols at one end or the other (or both). We let  $S_{4r}$  denote the labeling  $0_2 11 0_2 11 \dots 0_2 11$  (repeated  $r$ -times), Let  $S'_{4r}$  denote the labeling  $01_2 0 01_2 0 \dots 01_2 0$  (repeated  $r$ -times).

The labeling  $1_2 0_2 1_2 0_2 \dots 1_2 0_2$  (repeated  $r$ -times) and labeling  $10_2 110_2 1 \dots 10_2 1$  (repeated  $r$ -times) are written  $L_{4r}$  and  $L'_{4r}$ . Let  $M_r$  denote the labeling 01 01...01, zero-one repeated  $r$ times if  $r$  is even and 01 01...010 if  $r$  is

odd; for example,  $M_6 = 010101$  and  $M_5 = 01010$ . We let  $M'_r$  denote the labeling  $1010\dots10$ . Sometimes, we modify the labeling  $M_r$  or  $M'_r$  by adding symbols at one end or the other (or both). Also,  $L_{4r}$  (or  $L'_{4r}$ ) with extra labeling from right or left (or both sides) [9-12]. For specific labeling  $L$  and  $M$  of  $G \odot H$  where  $G$  is path and  $H$  is a second power of fan graph, we let  $[L; M]$  denote the corona labeling. Additional notation that we use is the following. For a given labeling of the corona  $G \odot H$ , we let  $v_i$  and  $e_i$  (for  $i = 0,1$ ) be the numbers of labels that are  $i$  as before, we let  $x_i$  and  $a_i$  be the corresponding quantities for  $G$ , and we let  $y_i$  and  $b_i$  be those for  $H$ , which are connected to the vertices labeled 0 of  $G$ . Likewise, let  $y'_i$  and  $b'_i$  be those for  $H$ , which are connected to the vertices labeled 1 of  $G$ . In case it increases by one more vertex, so  $y''_i$  and  $b''_i$  will be those for  $H$ , which are connected to the vertex labeled 1 or 0 of  $G$ . It is easy to verify that  $v_0 = x_0 + x_0y_0 + x_1y'_0, v_1 = x_1 + x_0y_1 + x_1y'_1, e_0 = a_0 + x_0b_0 + x_1b'_0 + x_0y_0 + x_1y'_1$  and  $e_1 = a_1 + x_0b_1 + x_1b'_1 + x_0(x_0y_1) + x_1y'_0$ . Thus  $|(v_0 + e_0) - (v_1 + e_1)| = |(x_0 - x_1) + (a_0 - a_1) + x_0(b_0 - b_1) + x_1(b'_0 - b'_1) + 2x_0(y_0 - y_1)|$ . Finally, for particular labeling A and B that are used for  $P_m$  and  $F_n^2$ .

### 3. The Total cordial of corona Product between paths and second power of Fan graphs

In this section, we explain that the corona between paths and the second power of Fan graphs  $P_m \odot F_n^2$  are Total cordial for all  $m \geq 1$ , and  $n \geq 4$ . This goal will be achieved after the following lemmas.

**Lemma 3.1**  $P_m \odot F_n^2$  is Total cordial for all  $m \geq 1$  and  $n \equiv 0(mod4)$ .

**Proof.** We should investigate the following cases :

**Case (1).  $m \equiv 0(mod4)$ .**

Let  $m = 4r, r \geq 1$ . Then, one can select the labeling  $[S_{4r}: 0M'_{4s}, 0 M'_{4s}, 1M_{4s}, 1M_{4s}, \dots (r - times)]$  for  $P_{4r} \odot F_{4s}^2$ . Therefore  $x_0 = x_1 = 2r, a_1 = 2r - 1, a_0 = 2r, y_0 = 2s + 1, y_1 = 2s, b_1 = 6s - 1, b_0 = 6s - 2, y'_0 = 2s, y'_1 = 2s + 1, b'_1 = 6s - 1$  and  $b'_0 = 6s - 2$ . Hence, It is simple to prove that  $|(v_0 + e_0) - (v_1 + e_1)| = 1$ . Thus  $P_{4r} \odot F_{4s}^2, s \geq 1$  is Total cordial. As an example, Figure (4.1) illustrates this case  $P_4 \odot F_4^2$ .

**Case (2).  $m \equiv 1(mod4)$ .**

Let  $m = 4r + 1, r > 0$ . Then, one can select the labelling  $[S_{4r+1}: 0M_{4s}, 0M_{4s}, 1M_{4s}, 1M_{4s}, \dots (r - times), 0M_{4s}]$  for  $P_{4r+1} \odot F_{4s}^2$ . Therefore  $x_0 = 2r, x_1 = a_1 = 2r - 1, a_0 = 2r + 1, y_0 = 2s + 1, y_1 = 2s, b_1 = 6s - 2, b_0 = 6s - 1, y'_0 = 2s, y'_1 = 2s + 1, b'_1 = 6s - 2, b'_0 = 6s - 1, y''_0 = 2s + 1, y''_1 = 2s, b''_1 = 6s - 2$  and  $b''_0 = 6s - 1$ . Hence, It is simple to prove that  $|(v_0 + e_0) - (v_1 + e_1)| = 0$ . Thus  $P_{4r+1} \odot F_{4s}^2, s \geq 1$  is Total cordial.

**Case (3).  $m \equiv 2(mod4)$ .**

Let  $m = 4r + 2, r > 0$ . Then, one can select the labelling  $[S_{4r+2}: 0M'_{4s}, 0M'_{4s}, 1M_{4s}, 1M_{4s}, \dots (r - times), 1M_{4s}, 0M'_{4s}]$  for  $P_{4r+2} \odot F_{4s}^2$ . Therefore  $x_0 = x_1 = 2r + 1, a_1 = 2r, a_0 = 2r + 1, y_0 = 2s + 1, y_1 = 2s, b_1 = 6s - 1, b_0 = 6s - 2, y'_0 = 2s, y'_1 = 2s + 1, b'_1 = 6s - 1$  and  $b'_0 = 6s - 2$ . Hence, It is simple to prove that  $|(v_0 + e_0) - (v_1 + e_1)| = 1$ . Thus  $P_{4r+2} \odot F_{4s}^2, s \geq 1$  is Total cordial.

**Case (4).  $m \equiv 3(mod4)$ .**

Let  $m = 4r + 3, r > 0$ . Then, one can select the labelling  $[S_{4r+3}: 0M'_{4s}, 0M'_{4s}, 1M'_{4s}, 1M'_{4s}, \dots (r - times), 1M'_{4s}, 1M_{4s}, 0M'_{4s}]$  for  $P_{4r+3} \odot F_{4s}^2$ . Therefore  $x_0 = 2r + 1, x_1 = a_1 = 2r, a_0 = 2r + 2, y_0 = 2s + 1, y_1 = 2s, b_1 = 6s - 1, b_0 = 6s - 2, y'_0 = 2s, y'_1 = 2s + 1, b'_1 = 6s - 1, b'_0 = 6s - 2, y''_0 = 2s + 1, y''_1 = 2s, b''_1 = 6s - 1$  and  $b''_0 = 6s - 2$ . Hence, It is simple to prove that  $|(v_0 + e_0) - (v_1 + e_1)| = 0$ . Thus  $P_{4r+3} \odot F_{4s}^2, s \geq 1$  is Total cordial.

**Lemma 3.2**  $P_m \odot F_n^2$  is Total cordial for all  $m \geq 1$  and  $n \equiv 1(mod4)$ .

**Proof.** We should investigate the following cases:

**Case (1).  $m \equiv 0(mod4)$ .**

Let  $m = 4r, r \geq 1$ . Then, one can select the labeling  $[S_{4r}: 11_3 0_2 M_{4s-4}, 11_3 0_2 M_{4s-4}, 0_4 1_2 M'_{4s-4}, 0_4 1_2 M'_{4s-4}, \dots (r - times)]$  for  $P_{4r} \odot F_{4s+1}^2$ . Therefore  $x_0 = x_1 = a_1 = 2r - 1, a_0 = 2r, y_0 = 2s, y_1 = 2s + 2, b_1 = 6s - 1, b_0 = 6s + 1, y'_0 = 2s + 2, y'_1 = 2s$  and  $b'_1 = 6s - 1, b'_0 = 6s + 1$ . Hence, It is simple to prove that  $|(v_0 + e_0) - (v_1 + e_1)| = 1$ . Thus  $P_{4r} \odot F_{4s+1}^2, s \geq 1$  is Total cordial. As an example, Figure (4.2) illustrates this case  $P_4 \odot F_5^2$ .

**Case (2).  $m \equiv 1(mod4)$ .**

Let  $m = 4r + 1, r > 0$ . Then, one can select the labeling  $[S_{4r}0: 11_3 0_2 M_{4s-4}, 11_3 0_2 M_{4s-4}, 0_4 1_2 M'_{4s-4}, 0_4 1_2 M'_{4s-4}, \dots (r - \text{times}), 10_3 1_2 M_{4s-4}]$  for  $P_{4r+1} \odot F_{4s+1}^2$ . Therefore  $x_0 = 2r + 1, x_1 = a_0 = a_1 = 2r, y_0 = 2s + 2, y_1 = 2s, b_1 = 6s + 1, b_0 = 6s - 1, y'_0 = 2s, y'_1 = 2s + 2, b'_1 = 6s + 1, b'_0 = 6s - 1, y''_0 = y''_1 = 2s + 1$ , and  $b''_0 = b''_1 = 6s$ . Hence, It is simple to prove that  $|(v_0 + e_0) - (v_1 + e_1)| = 1$ . Thus  $P_{4r+1} \odot F_{4s+1}^2, s \geq 1$ , is Total cordial.

**Case (3).  $m \equiv 2(\text{mod}4)$ .**

Let  $m = 4r + 2, r \geq 0$ . Then, one can select the labeling  $[S_{4r}101_4 0_2 M_{4s-4}, 11_3 0_2 M_{4s-4}, 0_4 1_2 M'_{4s-4}, 0_4 1_2 M'_{4s-4} \dots (r - \text{times}), 0_4 1_2 M'_{4s-4}, 11_3 0_2 M_{4s-4}]$  for  $P_{4r+2} \odot F_{4s+1}^2$ . Therefore  $x_0 = x_1 = a_1 = 2r, a_0 = 2r + 1, y_0 = 2s + 2, y_1 = 2s, b_1 = 6s + 1, b_0 = 6s - 1, y'_0 = 2s, y'_1 = 2s + 2$  and  $b'_1 = 6s + 1, b'_0 = 6s - 1$ . Hence It is simple to prove that  $|(v_0 + e_0) - (v_1 + e_1)| = 1$ . Thus  $P_{4r+2} \odot F_{4s+1}^2, s \geq 1$ , is Total cordial.

**Case (4).  $m \equiv 3(\text{mod}4)$ .**

Let  $m = 4r + 3, r \geq 0$ . Then, one can select the labeling  $[S_{4r}101: 1_4 0_2 M_{4s-4}, 1_4 0_2 M_{4s-4}, 0_4 1_2 M'_{4s-4}, 0_4 1_2 M'_{4s-4}, \dots (r - \text{times}), 0_4 1_2 M'_{4s-4}, 1_4 0_2 M_{4s-4}, 0_3 1_3 M'_{4s-4}]$  for  $P_{4r+3} \odot F_{4s+1}^2$ . Therefore  $x_0 = a_0 = a_1 = 2r + 1, x_1 = 2r + 2, y_0 = 2s, y_1 = 2s + 2, b_1 = 6s - 1, b_0 = 6s + 1, y'_0 = 2s + 2, y'_1 = 2s, b'_1 = 6s - 1, b'_0 = 6s + 1, y''_0 = y''_1 = 2s + 1$ , and  $b''_0 = b''_1 = 6s$ . Hence, It is simple to prove that  $|(v_0 + e_0) - (v_1 + e_1)| = 1$ . Thus  $P_{4r+3} \odot F_{4s+1}^2, s \geq 1$ , is Total cordial.

**Lemma 3.3**  $P_m \odot F_n^2$  is Total cordial for all  $m \geq 1$  and  $n \equiv 2(\text{mod}4)$ .

**Proof.** We need to study the following cases:

**Case (1).  $m \equiv 0(\text{mod}4)$ .**

Let  $m = 4r, r \geq 1$ . Then, one can select the labeling  $[S_{4r}: 0M'_{4s+2}, 0 M'_{4s+2}, 1M_{4s+2}, 1M_{4s+2}, \dots (r - \text{times})]$  for  $P_{4r} \odot F_{4s+2}^2$ . Therefore  $x_0 = x_1 = 2r, a_1 = 2r - 1, a_0 = 2r, y_0 = 2s + 2, y_1 = 2s + 1, b_1 = 6s + 1, b_0 = 6s + 2, y'_0 = 2s + 1, y'_1 = 2s + 2, b'_1 = 6s + 1$  and  $b'_0 = 6s + 2$ . Hence, It is simple to prove that  $|(v_0 + e_0) - (v_1 + e_1)| = 1$ . Thus  $P_{4r} \odot F_{4s+2}^2, r \geq 1$  is Total cordial. As an example, Figure (4.3) illustrates this case  $P_4 \odot F_6^2$ .

**Case (2).  $m \equiv 1(\text{mod}4)$ .**

Let  $m = 4r + 1, r \geq 0$ . Then, one can select the labeling  $[S_{4r}1: 0M'_{4s+2}, 0M'_{4s+2}, 1M'_{4s+2}, 1M'_{4s+2}, \dots, (r - \text{times}), 0M_{4s+2}]$  for  $P_{4r+1} \odot F_{4s+2}^2$ . Therefore  $x_0 = 2r, x_1 = 2r + 1, a_1 = 2r - 1, a_0 = 2r + 1, y_0 = 2s + 2, y_1 = 2s + 1, b_1 = 6s + 2, b_2 = 6s + 1, y'_0 = 2s + 1, y'_1 = 2s + 2, b'_1 = 6s + 2, b'_0 = 6s + 1, y''_0 = 2s + 2, y''_1 = 2s + 1, b''_1 = 6s + 2$  and  $b''_0 = 6s + 1$ . Hence, It is simple to prove that  $|(v_0 + e_0) - (v_1 + e_1)| = 0$ . Thus  $P_{4r+1} \odot F_{4s+2}^2, s \geq 1$  is Total cordial.

**Case (3).  $m \equiv 2(\text{mod}4)$ .**

Let  $m = 4r + 2, r \geq 0$ . Then, one can select the labeling  $[S_{4r}10: 0 M'_{4s+2}, 0M'_{4s+2}, 1M_{4s+2}, 1M_{4s+2}, \dots, (r - \text{times}), 1M_{4s+2}, 0M'_{4s+2}]$  for  $P_{4r+2} \odot F_{4s+2}^2$ . Therefore  $x_0 = x_1 = a_0 = 2r + 1, a_1 = 2r, y_0 = 2s + 2, y_1 = 2s + 1, b_1 = 6s + 2, b_0 = 6s + 1, y'_0 = 2s + 1, y'_1 = 2s + 2, b'_1 = 6s + 2$  and  $b'_0 = 6s + 1$ . Hence, It is simple to prove that  $|(v_0 + e_0) - (v_1 + e_1)| = 1$ . Thus  $P_{4r+2} \odot F_{4s+2}^2, s \geq 1$  is Total cordial.

**Case (4).  $m \equiv 3(\text{mod}4)$ .**

Let  $m = 4r + 3, r \geq 0$ . Then, one can select the labeling  $[S_{4r}1_2 0: 0M'_{4s+2}, 0M'_{4s+2}, 1M_{4s+2}, 1M_{4s+2}, \dots, (r - \text{times}), 1M_{4s+2}, 0_3 1_2 M'_{4s-2}, 0M'_{4s+2}]$  for  $P_{4r+3} \odot F_{4s+2}^2$ . Therefore  $x_0 = 2r + 1, x_1 = a_0 = 2r + 2, a_1 = 2r, y_0 = 2s + 2, y_1 = 2s + 1, b_1 = 6s + 2, b_0 = 6s + 1, y'_0 = 2s + 1, y'_1 = 2s + 2, b'_1 = 6s + 2, b'_0 = 6s + 1, y''_0 = 2s + 2, y''_1 = 2s + 1, b''_1 = 6s + 2$  and  $b''_0 = 6s + 1$ . Hence, It is simple to prove that  $|(v_0 + e_0) - (v_1 + e_1)| = 0$ . Thus  $P_{4r+3} \odot F_{4s+2}^2, s \geq 1$  is Total cordial.

**Lemma 3.4**  $P_m \odot F_n^2$  is Total cordial for all  $m \geq 1$  and  $n \equiv 3(\text{mod}4)$ .

**Proof:** We should examine the following cases:

**Case (1).  $m \equiv 0(\text{mod}4)$ .**

Let  $m = 4r, r \geq 1$ . Then, one can select the labeling  $[S_{4r}: 10_3 M'_{4s}, 10_3 M'_{4s}, 101_2 M_{4s}, 101_2 M_{4s}, \dots, (r - \text{times})]$  for  $P_{4r} \odot F_{4s+3}^2$ . Therefore  $x_0 = x_1 = a_0 = 2r, a_1 = 2r - 1, y_0 = 2s + 3, y_1 = 2s + 1, b_1 = 6s + 4, b_0 = 6s +$

$2, y'_0 = 2s + 1, y'_1 = 2s + 3$  and  $b'_1 = 6s + 4, b'_0 = 6s + 2$ . Hence, It is simple to prove that  $|(v_0 + e_0) - (v_1 + e_1)| = 1$ . Thus  $P_{4r} \odot F_{4s+3}^2, s \geq 1$  is Total cordial. As an example, Figure (4.4) illustrates this case  $P_4 \odot F_7^2$ .

**Case (2).  $m \equiv 1 \pmod{4}$ .**

Let  $m = 4r + 1, r \geq 0$ . Then, one can select the labeling  $[S_{4r+1}: 10_3M'_{4s}, 10_3M'_{4s}, 101_2M_{4s}, 101_2M_{4s}, \dots, (r - \text{times}), 01_20M'_{4s}]$  for  $P_{4r+1} \odot F_{4s+3}^2$ . Therefore  $x_0 = 2r, x_1 = a_1 = 2r - 1, a_0 = 2r + 1, y_0 = 2s + 3, y_1 = 2s + 1, b_1 = 6s + 2, b_0 = 6s + 4, y'_0 = 2s + 1, y'_1 = 2s + 3, b'_1 = 6s + 2, b'_0 = 6s + 4, y''_0 = y''_1 = 2s + 2, b''_1 = 6s + 2$  and  $b''_0 = 6s + 4$ . Hence It is simple to prove that  $|(v_0 + e_0) - (v_1 + e_1)| = 1$ . Thus  $P_{4r+1} \odot F_{4s+3}^2, s \geq 1$  is Total cordial.

**Case (3).  $m \equiv 2 \pmod{4}$ .**

Let  $m = 4r + 2, r \geq 0$ . Then, one can select the labeling  $[S_{4r+2}: 10_3M'_{4s}, 10_3M'_{4s}, 101_2M_{4s}, 101_2M_{4s}, \dots, (r - \text{times}), 101_2M_{4s}, 10_3M'_{4s}]$  for  $P_{4r+2} \odot F_{4s+3}^2$ . Therefore  $x_0 = x_1 = a_1 = 2r + 1, a_0 = 2r, y_0 = 2s + 3, y_1 = 2s + 1, b_1 = 6s + 4, b_0 = 6s + 2, y'_0 = 2s + 1, y'_1 = 2s + 3, b'_1 = 6s + 4$  and  $b'_0 = 6s + 2$ . Hence, It is simple to prove that  $|(v_0 + e_0) - (v_1 + e_1)| = 1$ . Thus  $P_{4r+2} \odot F_{4s+3}^2, s \geq 1$  is Total cordial.

**Case (4).  $m \equiv 3 \pmod{4}$ .**


Let  $m = 4r + 3, r \geq 0$ . Then, one can select the labeling  $[S_{4r+3}: 10_3M'_{4s}, 10_3M'_{4s}, 101_2M_{4s}, 101_2M_{4s}, \dots, (r - \text{times}), 101_2M_{4s}, 10_3M'_{4s}, 01_20M'_{4s}]$  for  $P_{4r+3} \odot F_{4s+3}^2$ . Therefore  $x_0 = a_0 = 2r + 2, x_1 = 2r + 1, a_1 = 2r, y_0 = 2s + 3, y_1 = 2s + 1, b_1 = 6s + 4, b_0 = 6s + 2, y'_0 = 2s + 1, y'_1 = 2s + 3, b'_1 = 6s + 4, b'_0 = 6s + 2, y''_0 = y''_1 = 2s + 2, b''_1 = 6s + 4$  and  $b''_0 = 6s + 2$ . Hence It is simple to prove that  $|(v_0 + e_0) - (v_1 + e_1)| = 1$ . Thus  $P_{4r+3} \odot F_{4s+3}^2, s \geq 1$  is Total cordial.

The following theorem may be established as a result of all preceding lemmas.


**Theorem 3.1.** The corona between path and fourth power of fan graphs  $P_m \odot F_n^2$  is Total cordial for all  $m \geq 1$  and  $n \geq 4$ .

**4. Example**

The Total cordial graph of  $P_4 \odot F_4^2, P_4 \odot F_5^2, P_4 \odot F_6^2$  and  $P_4 \odot F_7^2$  are illustrated in Figures (4.1, ..., 4.4).




**Figure (4.1).**  $P_4 \odot F_4^2$  is Total cordial




$$|(v_0 + e_0) - (v_1 + e_1)| = 1$$

Figure (4.2).  $P_4 \odot F_5^2$  is Total cordial.



$$|(v_0 + e_0) - (v_1 + e_1)| = 1$$

Figure (4.3).  $P_4 \odot F_6^2$  is Total cordial



$$|(v_0 + e_0) - (v_1 + e_1)| = 1$$

Figure (4.4).  $P_4 \odot F_7^2$  is Total cordial.

### 5. Conclusion

In this work, we proved that the corona between paths and second power of Fan graphs  $P_m \odot F_n^2$  is Total cordial for all  $m \geq 1$ , and  $n \geq 4$ . An example is introduced in section 4.

### References

- [1] Cahit, I. (1987). Cordial graphs-a weaker version of graceful and harmonious graphs. *Ars combinatoria*, 23, 201-207.
- [2] Gallian, J. A. (2012). Graph labeling. *The electronic journal of combinatorics*, DS6-Dec
- [3] Golomb, S. W. (1972). How to number a graph. In *Graph theory and computing* (pp. 23-37). Academic press.
- [4] Graham, R. L., Sloane, N. J. A. (1980). On additive bases and harmonious graphs. *SIAM Journal on Algebraic Discrete Methods*, 1(4), 382-404.
- [5] Harary, F. and Wilcox, G. W., Boolean operations on graphs, *Math.Scand.*20, pp.41-51(1967).

- [6] Sundaram, M., Ponraj, P. and Somasundaram, S., "Product cordial labeling of graphs", *Bulletin of Pure and Applied Sciences*, 23E (1) (2004), 155-162.
- [7] Sundaram, M., Ponraj, P. and Somasundaram, S., "Total Product Cordial Labeling of Graphs", *Bulletin of Pure and Applied Sciences*, 23E (1) (2005), 199-203.
- [8] Aldous, J. M., Wilson, R. J. (2003). *Graphs and applications: an introductory approach*. Springer Science and Business Media.
- [9] Harray, F. (1953). On the notion of a signed graph. *Michigan Math. J*, 2, 143-146.
- [10] Hovey, M. (1991). A-cordial graphs. *Discrete Mathematics*, 93(2-3), 183-194.
- [11] Kirchherr, W. W. (1991). On the cordiality of some specific graphs. *Ars Combinatoria*, 31, 127-138.
- [12] Lee, S. M., Liu, A. (1991). A construction of cordial graphs from smaller cordial graphs. *Ars Combinatoria*, 32, 209-214.
- [13] Rosa, A. (1966, July). On certain valuations of the vertices of a graph. In *Theory of graphs (Internat. symposium, Rome)* (pp. 349-355).
- [14] Seoud, M. A., Diab, A. T., Elsayhawi, E. A. (1998). On strongly-C harmonious, relatively prime, odd graceful and cordial graphs. In *Proc. Math. Phys. Soc. Egypt (Vol. 73)*, pp. 33-55).
- [15] ELrokh, A., Ali Al-Shamiri, M. M., Nada, S., & El-hay, A. A. (2022). Cordial and Total Cordial Labeling of Corona Product of Paths and Second Order of Lemniscate Graphs. *Journal of Mathematics*, 2022.
- [16] Badr, E., Nada, S., Ali Al-Shamiri, M. M., Abdel-Hay, A., & ELrokh, A. (2022). A novel mathematical model for radio mean square labeling problem. *Journal of Mathematics*, 2022.
- [17] ELrokh, A., Ali Al-Shamiri, M. M., & El-hay, A. (2022). A Novel Problem to Solve the Logically Labeling of Corona between Paths and Cycles. *Journal of Mathematics*, 2022.
- [18] Badr, E., Abd El-hay, A., Ahmed, H., & Moussa, M. (2021). Polynomial, exponential and approximate algorithms for metric dimension problem. *Mathematical Combinatorics*, 2, 51-67.