

# SOLVING A BIRKHOFF INTERPOLATION PROBLEM FOR (0,1,5) DATA

Rama Nand Mishra<sup>a</sup>, Akhilesh Kumar Mishra<sup>b</sup>, and Kulbhushan Singh<sup>c</sup>

<sup>a</sup>Research scholar , Faculty of Mathematical and Statistical Sciences, Sri Ramswaroop Memorial University, Lucknow-Deva Road, Barabanki, U.P., India, [ramanandmishra.ids@srmu.ac.in](mailto:ramanandmishra.ids@srmu.ac.in)

<sup>b</sup>Associate Professor, Faculty of Mathematical and Statistical Sciences, Sri Ramswaroop Memorial University, Lucknow-Deva Road, Barabanki, U.P., India, [akhileshmishra.maths@srmu.ac.in](mailto:akhileshmishra.maths@srmu.ac.in)

<sup>c</sup>Professor, Faculty of Mathematical and Statistical Sciences , Sri Ramswaroop Memorial University, Lucknow-Deva Road, Barabanki, U.P., India , [kulbhushan.maths@srmu.ac.in](mailto:kulbhushan.maths@srmu.ac.in)

## ABSTRACT

The following research work deals with a special type of Birkhoff interpolation problem in which we have 3 set of data prescribed in the unit interval  $I = [0, 1]$ . Data values are the function value, first derivatives and fifth derivatives prescribed at nodes of the unit interval. We obtained a unique spline interpolating the given data along with the convergence problem.

**Keywords:** Spline function, Lacunary Interpolation, Birkhoff interpolation.

## 1. INTRODUCTION

In this study, we use virtually quantic splines  $s(x) \in S_{n,5}^{(2)}(x)$  for a given partition  $\Delta$  to solve a Birkhoff interpolation issue we term (0, 1, 5) problem.

Let  $\Delta : 0 = x_0, x_1, \dots, x_{n-1}, x_n = 1$  be a partition of the unit interval  $I = [0,1]$  and  $S_{n,5}^{(2)}(x)$  denotes the class of spline function  $s(x)$  such that

$$(1.1) \quad \begin{aligned} s_i(x) &\in \pi_5, & i &= 0(1)n-2 \\ s_i(x) &\in \pi_6, & i &= 0(1)n-1 \quad \text{where } x \in [x_i, x_{i+1}] \end{aligned}$$

$$(1.2) \quad s(x) \in C^2(I).$$

Also, we denote  $x_{i+1} - x_i = h_i$ , for all  $i = 0(1)n - 1$ . We prove the existence and uniqueness of such spline functions and show that they converge to the given function  $f(x) \in C^5(I)$  up to derivative of order 5. For relevant reading one is referred to [1 -7].

Following two theorems were proved:

## 2. THEOREM OF UNIQUE EXISTENCE

Given  $\Delta$  and the real numbers  $y_i, y'_i, y_i^{(5)}, i = 0(1)n, y_0'', y_n''$  there exists unique  $s_\Delta(x) \in S_{n,5}^{(2)}(x)$  such that

$$(2.1) \quad s_\Delta^{(q)}(x_i) = y_i^{(q)}, \quad i = 0(1)n - 2 \quad q = 0,1,5$$

$$(2.2) \quad s_\Delta''(x_0) = y_0'', \quad s_\Delta''(x_n) = y_n''$$

**Proof :**

We set

$$(2.3) \left\{ \begin{array}{l} s_0(x), \text{ where } x \in [x_0, x_1] \\ s_\Delta(x) = s_i(x), \text{ when } x \in [x_i, x_{i+1}] \text{ } i=0(1) n-2. \\ s_{n-1}(x), \text{ when } x \in [x_{n-1}, x_n] \end{array} \right.$$

Using interpolatory conditions we write,

$$(2.4) \quad s_0(x) = y_0 + \frac{(x-x_0)}{1!}y_0' + \frac{(x-x_0)^2}{2!}y_0'' + \frac{(x-x_0)^3}{3!}a_{0,3} + \frac{(x-x_0)^4}{4!}a_{0,4} + \frac{(x-x_0)^5}{5!}y_0^{(5)} + \frac{(x-x_0)^6}{6!}a_{0,6}$$

$$(2.5) \quad s_i(x) = y_i + \frac{(x-x_i)}{1!}y_i' + \frac{(x-x_i)^2}{2!}a_{i,2} + \frac{(x-x_i)^3}{3!}a_{i,3} + \frac{(x-x_i)^4}{4!}a_{i,4} + \frac{(x-x_i)^5}{5!}y_i^{(5)}$$

$$(2.6) \quad s_{n-1}(x) = y_{n-1} + \frac{(x-x_{n-1})}{1!}y_{n-1}' + \frac{(x-x_{n-1})^2}{2!}a_{n-1,2} + \frac{(x-x_{n-1})^3}{3!}a_{n-1,3} + \frac{(x-x_{n-1})^4}{4!}a_{n-1,4} + \frac{(x-x_{n-1})^5}{5!}y_{n-1}^{(5)} + \frac{(x-x_{n-1})^6}{6!}a_{n-1,6}$$

The coefficients involved in the above equation are determined by the remaining interpolatory conditions and the continuity requirement that  $s_\Delta(n) \in C^2(I)$ . Applying these condition we get the following set of equations.

$$(2.7) \left\{ \begin{array}{l} y_1 = y_0 + h_0y_0' + \frac{h_0^2}{2!}y_0'' + \frac{h_0^3}{3!}a_{0,3} + \frac{h_0^4}{4!}a_{0,4} + \frac{h_0^5}{5!}y_0^{(5)} + \frac{h_0^6}{6!}a_{0,6} \\ y_1' = y_0' + h_0y_0'' + \frac{h_0^2}{2!}a_{0,3} + \frac{h_0^3}{3!}a_{0,4} + \frac{h_0^4}{4!}y_0^{(5)} + \frac{h_0^5}{5!}a_{0,6}, \\ y_1^{(5)} = y_0^{(5)} + h_0a_{0,6} \end{array} \right.$$

$$(2.8) \left\{ \begin{array}{l} y_{i+1} = y_i + h_iy_i' + \frac{h_i^2}{2!}a_{i,2} + \frac{h_i^3}{3!}a_{i,3} + \frac{h_i^4}{4!}a_{i,4} + \frac{h_i^5}{5!}y_i^{(5)} \\ y_{i+1}' = y_i' + h_ia_{i,2} + \frac{h_i^2}{2!}a_{i,3} + \frac{h_i^3}{3!}a_{i,4} + \frac{h_i^4}{4!}y_i^{(5)} \\ y_{i+1}^{(5)} = y_i^{(5)} \end{array} \right.$$

$$(2.9) \left\{ \begin{array}{l} y_n = y_{n-1} + h_{n-1}y_{n-1}' + \frac{h_{n-1}^2}{2!}a_{n-1,2} + \frac{h_{n-1}^3}{3!}a_{n-1,3} + \frac{h_{n-1}^4}{4!}a_{n-1,4} \\ \quad + \frac{h_{n-1}^5}{5!}y_{n-1}^{(5)} + \frac{h_{n-1}^6}{6!}a_{n-1,6} \\ y_n' = y_{n-1}' + \frac{h_{n-1}}{1!}a_{n-1,2} + \frac{h_{n-1}^2}{2!}a_{n-1,3} + \frac{h_{n-1}^3}{3!}a_{n-1,4} + \frac{h_{n-1}^4}{4!}y_{n-1}^{(5)} \\ \quad + \frac{h_{n-1}^5}{5!}a_{n-1,6} \\ y_n^{(5)} = y_{n-1}^{(5)} + h_{n-1,6} \end{array} \right.$$

From these equations, we have

$$(2.10) \left\{ \begin{array}{l} a_{0,3} = 6h_0^{-3}(4y_1 - 4y_0 - h_0y_1' - 3h_0'y_0' - h_0^2y_0'') \\ \quad + \frac{1}{60}h_0^2(2y_0^{(5)} + y_1^{(5)}) \\ a_{0,4} = 24h_0^{-3}(3h_0^{-1} - 3h_0^{-1}y_1 + y_1' + 2y_0') \\ \quad + 12h_0^{-2}y_0'' - \frac{1}{10}h_0(6y_0^{(5)} - y_1^{(5)}) \\ a_{0,6} = h_0^{-1}(h_1^{(5)} - y_0^{(5)}) \\ \quad \vdots \\ \quad \vdots \end{array} \right.$$

$$(2.11) \left\{ \begin{aligned} a_{i,2} &= h_i^{-2}(2y_i - h_i y'_{i+1} - 16h_i y'_i - 4h_i^2 y''_i) \\ &\quad + \frac{7}{180} h_i^3 y_i^{(5)} - \frac{1}{72} h_i^3 y_{i+1}^{(5)}, \\ a_{i,3} &= 6h_i^{-3}(4y_{i+1} - h_i y'_{i+1} - 4y_i + 5h_i y'_i - h_i^2 y''_i) \\ &\quad + \frac{1}{20} h_i^2 (3y_i^{(5)} + y_{i+1}^{(5)}), \\ a_{i,4} &= 24h_i^{-3} \left( -3h_i^{-1} y_{i+1} + \frac{3}{2} h_i^{-1} y'_{i+1} + 3h_i^{-1} y_i + 2h_i y'_i + h_i y''_i \right) \\ &\quad - \frac{h_i}{30} (11y_i^{(5)} - y_{i+1}^{(5)}), \end{aligned} \right.$$

And

$$(2.12) \left\{ \begin{aligned} a_{n-1,2} &= 3y_{n-1}'' + \frac{h_{n-1}^3}{360} (2083y_{n-1}^{(5)} - 649y_n^{(5)}), \\ a_{n-1,3} &= 6h_{n-1}^{-3} (4y_n - h_{n-1} y'_n - 4y_{n-1} - 3h_{n-1} y'_{n-1} - h_{n-1}^2 y''_{n-1}) \\ &\quad + \frac{1}{60} h_{n-1}^2 (11y_{n-1}^{(5)} + y_n^{(5)}), \\ a_{n-1,4} &= 24h_{n-1}^{-4} (-3y_n + h_{n-1} y'_n + 3y_{n-1} + 2h_{n-1} y'_{n-1} + \frac{1}{2} h_{n-1}^2 y''_{n-1}) \\ &\quad - \frac{1}{30} h_{n-1} (5y_{n-1}^{(5)} - y_n^{(5)}) \\ a_{n-1,6} &= h_{n-1}^{-1} (y_n^{(5)} - y_{n-1}^{(5)}) \end{aligned} \right.$$

The unique existence of the coefficients above show the unique existence of the spline function  $s_{\Delta}(xx)$  of theorem 1. This is the prove of the theorem.

### 3. THEOREM OF CONVERGENCE

Let  $ff(xx) \in CC^5(1)$ , then the unique spline function  $ss_{\Delta}(xx)$  mentioned in theorem 1, with  $yy_{ii}$ , etc being associated with the function  $ff(xx)$ ,  $iee yy_{ii} = ff(xx_i)$ ,  $yy_{ii} = ff'(xx_i)$  etc., we have for  $xx \in [xx_i, xx_{i+1}]$ ,  $i = 0(1)nn - 1$ ,

$$(3.1) |ss_{\Delta}^{(qq)}(xx) - ff^{(qq)}(xx)| \leq kkqq h^{5-qq} ww_5(h), qq = 0(1)5$$

Here we take  $h_i = h$  for all  $i = 0(1)n-1$  and denote the modulus of continuity of  $ff(xx) \in CC^5(II)$  by  $w_5(h)$ .

**Proof :**

Let first be  $xx_{ii} \leq xx \leq xx_{ii+1}$ ,  $ii = 0(1)nn - 2$ , from (2.4) and writing finite Taylor's sums for  $f(xx)$  and its derivatives, we have

$$s_i^{(q)}(x) - f^{(q)}(x) = \frac{(x - x_i)^{5-q}}{(5 - q)!} (a_{i,5} - f^{(5)}(x_i)) + \frac{(x - x_i)^{6-q}}{(6 - q)!} (a_{i,6} - f^{(6)}(x_i))$$

When  $q = 4, 5$ ; where  $x_i < \xi_{i,q} < x_{i+1}$ .

Let  $x \in [x_0, x_1]$ , then  $q = 5, i = 0(1)n$

$$\begin{aligned} s_0^{(5)}(x) - f^{(5)}(x) &= (x - x_0)(a_{i,6} - f^{(6)}(x)) \\ &= (x - x_0)[h_0^{-1}(y_1^{(5)} - y_0^{(5)}) - f^{(6)}(x)] \end{aligned}$$

Therefore,

$$|s_0^{(5)}(x) - f^{(5)}(x)| \leq w_5(h)$$

Using the interpolatory conditions, we have

$$|s_0^{(4)}(x) - f^{(4)}(x)| = \left| \int_{x_0}^{x_1} [s_0^{(5)}(x) - f^{(5)}(x)] dx \right|$$

$$\leq |x_1 - x_0| |s_0^{(5)}(x) - f^{(5)}(x)|$$

$$\leq h w_5(h)$$

Again using Taylor's theorem, we have

$$|s_0''(x) - f''(x)| \leq h^2 w_5(h)$$

Further,

$$\begin{aligned} |s_0''(x) - f''(x)| &\leq \left| \int_{x_0}^{x_1} \{s_0''(x) - f''(x)\} dx \right| \\ &\leq |x_1 - x_0| |s_0''(x) - f''(x)| \\ &\leq h^3 w_5(h) \end{aligned}$$

Similar

$$|s_0'(x) - f'(x)| \leq h^4 w_5(h)$$

$$\text{And } |s_0(x) - f(x)| \leq h^5 w_5(h).$$

This is the prove of theorem 2 for  $x \in [x_0, x_1]$ .

For  $x \in [x_i, x_{i+1}]$ ,  $i = (1)n - 2$ , we have (2.5) using Taylor's theorem

$$\begin{aligned} |s_i^{(5)}(x) - f^{(5)}(x)| &= |f_{i,5}^{(5)}(x) - f^{(5)}(x)| \\ &= |ff_{i,5}^{(5)}(nn) - ff_{i,5}^{(5)}(xx) + ff_{i,5}^{(5)} - ff^{(5)}(xx)| \\ &\leq ww_5(h)(wwheerree yy_{ii+1}^{(5)} = yy_{ii}^{(5)}.) \end{aligned}$$

This gives the results for  $xx \in [xx_{ii}, xx_{ii+1}]$ ,  $ii = (1)nn - 2$ . Proof for  $xx \in [xx_{nn-1}, xx_{nn}]$  can be carried out on

similar lines, so we omit the details.

#### 4. CONCLUSION

We have taken here a (0,1,5) lacunary interpolation problem for which we found a quantic spline function.  $ss(xx) \in SS_{nn,5}^{(2)}(xx)$  which interpolates the given data. Also it is shown that this spline function converges uniquely by finding error bounds. Such types of spline function can be used for solving differential equations and obtaining quatratre formula.

#### 5. REFERENCES

- [1]. Loscalzo, F.R., Talbot, T. D., (1967) Spline and approximation for solutions of ordinary differential equations. SIAM J. Numer. Anal., 4: 433-445.
- [2]. Micula, Gh., (1973) Approximate solution of the differential equation  $y''(x) = f(x,y)$  with spline functions. Math. ofcomput, 27: 807-816.
- [3]. Fawzy, T. (1977) Spline functions and the Cauchy's problem II. Acta Math. Hung, 29 (3-4): 259-271.
- [4]. Gyovari, J., (1984) Lakunare spline funktionun das Cauchy problem, Acta Math Hung, 44 (3-4): 327-335.
- [5]. Sallam, S., Hussain, M. A., (1984) deficient spline for approximation to second order differential equations. Appl. Math Modeling, 8: 408-412.
- [6]. Singh, K. B., Pandey, Ambrish Kumar., Ahmad, QaziShoeb., (2012), Solution of a Birkhoff Interpolation Problem by a Special Spline Function. International J. of Comp. App., 48: 22-27,
- [7]. Pandey, A. K., Ahmad, Q. S., Singh, K., (2013) Lacunary Interpolation (0,2;3) Problem and Some Comparison from Quartic Splines. American J. of App. Math. And Statistics, 1(6): 117-120.