

## Associated Factors Of Metabolic Syndrome And The Role Of Oral Health Among 50 And Older Adults: from the 6th KNHANES

Moon-Hee Kwon

Dept. of Nursing, Kyungdong University, Wonju-city Gangwon-do, 26495, KOREA

**Article History:** Received: 11 november 2020; Accepted: 27 December 2020; Published online: 05 April 2021

**Abstract:** The study on relationship between metabolic syndrome and oral health, is rare. We investigated to find out the correlation of metabolic syndrome and oral health. We investigated to find out the correlation of metabolic- syndrome and oral health with the data from the 6th Korea- National- Health and Nutrition- Examination- Survey. This is the cross-sectional designed study. 2433 subjects (aged 50 ≤) were included. Metabolic- syndrome was defined as NCEP-ATP (National- Cholesterol- Education- Program-Adult- Treatment- Panel III). To analyze influencing factors of metabolic syndrome, multi-variable- logistic- regression was used. Our results showed that gender(male) (OR 1.86; 95%CI: 1.27-2.72; p<.05) and not taking dietary supplements (OR 1.74; 95%CI: 1.22-2.48; p<.05) factors were the most influencing factors of metabolic syndrome. And then, presence of stress(OR1.59; 95%CI: 1.28-1.98; p<.05), lowest education level(elementary)(OR1.48; 95%CI: 0.98-2.21; p=.05), and oral health(periodontitis) (OR1.39; 95%CI: 1.09-1.78; p<.05) factors were showed as the next strong factors of metabolic syndrome in order. While chewing ability was not related with metabolic syndrome. We highlighted on the impact of oral health in metabolic syndrome including chewing difficulty as well as periodontitis. And to prevent metabolic syndrome cause of serious disability and lowering quality of life, we tried to find out the influencing factors of metabolic syndrome considering of this study with diverse view point such as adding the dietary supplements as dietary habit, stress as psychological value, and sedentary time as health-related factors. The government and public health associates should do urgently preparing the policies and strategies focusing on gender disparities, dietary supplements and oral health to prevent metabolic syndrome among ageing population.

**Keywords:** Dietary supplements, Education level, Gender, Metabolic syndrome, Periodontitis, Stress.

### Introduction

Metabolic syndrome was prevalent and increasing worldwide. Metabolic syndrome, the serious health problem resulted in cardiovascular disease, consists of hypertension, glucose intolerance, abdominal obesity, hypertriglyceridemia and low HDL cholesterol. Recently oral health factors, the major indicator of quality of life (Cervino, G., et al, 2019), were reported to being associated with metabolic- syndrome (Kim, J. S., et al, 2019; Kaur, K. K. et al, 2019). And gender disparities were also reported to be related with metabolic syndrome (Kim, J. S., et al, 2019) as well as the relationship between metabolic syndrome and healthy life style factors (Hoyas, I., &Leon-Sanz, M. 2019; Lin, W. T., et al, 2019). And coping style and occupational stress were reported to be associated with periodontitis among workers by the cross-sectional study(Islam, M. M., et al, 2019). Also, circulation microparticles in metabolic syndrome with cardiovascular risks were reported to be associated with oxidative stress (Zahran, A. M., et al, 2019). And posttraumatic stress was reported to be related with metabolic syndrome especially in women(Lihua, M., et al, 2020).In current evidence on natural agents, they were reported to be on thoughtful perspective in oral health including periodontitis (Jaramillo Flores, M. E. ,2019). It was reported on the links that serum vitamin C and antioxidant levels were related with periodontitis (Isola, G., et al, 2019; Van der Veldena, U., 2020). Especially, the relationship between vitamin D level and metabolic syndrome was suggested as important one in obese patient as well as general people for reducing metabolic disease and increasing longevity (Moukayed, M., & Grant, W. B.,2019). And dietary nitrate was also reported to be associated with activating of metabolic pathways and reducing of oxidative stress (Lundberg, J. O.,2018). Dietary supplements such as botanical oil and fish oil was reported to be related with metabolic syndrome as well as type-2 diabetes ( Lee, T. C. et al, 2014). In addition to that, dietary supplementation of watermelon juice was reported to being ameliorating the metabolic syndromes in obese animals (Wu, G., Collins, et al, 2007) though it has kinds of some limitation in animal subjects. And more, oral health was reported to be closely connected with improving of overall health especially among adults (Onubogu, U., et al, 2019; Ismail, F., et al, 2019; Rawal, I., et al, 2019). In addition to that, number of teeth, as the important indicator of oral health, was reported to increase the risk of metabolic syndrome among elderly (PEDRO, R. E. L., et al, 2019). Additionally, diabetes and smoking were closely associated with poor oral health such as tooth loss by recent cohort study( Kim, Y. T., et al, 2019). Through reviewing of current studies, we investigated to find out the major influencing factors of metabolic syndrome and impacting factors of oral-health of 50 and older aged Koreans using nationally representative data of 6th KNHANES. Considering potential impact on subjects and lifestyle, we included demographic, socio-economic, health related and oral health factors.

### Materials and Methods

\*Corresponding author: Moon-Hee Kwon

Dept. of Nursing, Kyungdong University, Wonju-city Gangwon-do, 26495, KOREA

Using derived from nationally representative data from 6th Knhanes (Korea- National- Health and Nutrition- Examination- Survey), of stratified sampling one, after exclusion of missing and not available one, this study was analyzed. The standards of NCEP-ATPⅢ (National- Cholesterol- Education- Program -Adult- Treatment- Panel) for the criteria of metabolic- syndrome was used for the study. This study is a cross-sectional- designed one.

### Study-subjects

This study was conducted to find-out on relationship between oral health and metabolic- syndrome of 2433 consented participants, 50 aged and older, using complex-sampling-methods by multiple-logistic-regressions with SPSS (ver. 21.0).

### Study-analysis

This study was analyzed by complex sampling methods. For the general characteristic of the 2433 subjects of aged 50 and over, demographic statistics methods were used. And chi-square test was used for analyzing prevalence of association between metabolic- syndrome and determinant factors. To investigate influencing factors of metabolic- syndrome and impact of oral health, multi-variable logistic regression methods with SPSS (ver. 21.0) were used. The significance- level was ( $p < .05$ ).

### Results and Discussion

Out of 2433 participants, 42.0% were men [Table 1]. 50-64 aged group was 52.2%. The prevalence of metabolic syndrome was 13.6%. About 30.2% participants reported no-taking of dietary supplements. 73.2% participants had stress. The lowest educational group(elementary) was 42.9%. Around 38.1% participants had periodontitis.

**Table 1. General characteristics of 50 and older participants**

Variables	N (2433)	%
Gender		
Male	1023	42.0
Female	1410	58.0
Age		
50-64	1271	52.2
65 over	1162	47.8
Residence		
Urban (Dong)	1851	76.1
Rural (Eup, Myeon)	582	23.9
Marital status(spouse)		
No	27	1.1
Yes	2406	98.9
Income level		
Lowest	562	23.1
Middle-low	625	25.7
Middle-high	615	25.3
Highest	631	25.9
Education level		
Elementary	1044	42.9
Middle-school	428	17.6
High-school	593	24.4
College	368	15.1
Smoking		
No	1506	61.9
Yes	927	38.1
Drinking		
No	480	19.7
Yes	1953	80.3

Physical activity(exercise)			
	No	1860	76.4
	Yes	573	23.6
Stress recognition			
	No	653	26.8
	Yes	1780	73.2
Taking dietary supplements			
	No	735	30.2
	Yes	1698	69.8
Sedentary time			
	Less than 5 hrs.	662	27.2
	Over 5 hrs.	1771	72.8
Chewing difficulty			
	Severe	919	37.8
	Moderate	405	16.6
	None	1109	45.6
Periodontitis			
	No	1506	61.9
	Yes	927	38.1
Metabolic syndrome			
	No	2101	86.4
	Yes	332	13.6

Table 2. show prevalence of association between metabolic- syndrome and determinant- factors with impact of oral health among Korean 50 and older adults. The prevalence of metabolic- syndrome was 55.7% (men) and 44.3% (women). And prevalence of metabolic syndrome was showed as smoking (50.3%), yes feeling stress (64.5%), not taking dietary supplements (35.5%), and periodontitis (45.5%).

**Table 2. Prevalence of Association between metabolic syndrome and determinant factors**

	Metabolic syndrome		P-value	
	Yes	No		
	N(%)	N(%)		
Variables				
Gender			<.001	
	Male	185(55.7)		838(39.9)
	Female	147(44.3)		1263(60.1)
Age			.137	
	50-64	186(56.0)		1085(51.6)
	65 over	146(44.0)		1061(48.4)
Residence			.374	
	Urban(Dong)	2509(78.0)		1592(75.8)
	Rural(Eup, Myeon)	73(22.0)		50.9(24.2)
Marital status(spouse)			.458	
	No	5(1.5)		22(1.0)
	Yes	327(98.5)		2079(99.0)
Income level			.895	
	Lowest	78(23.5)		484(23.0)
	Middle-low	85(25.6)		540(25.7)

Middle-high	88(26.5)	527(25.1)	
Highest	81(24.4)	550(26.2)	
Education level			.897
Elementary	142(42.8)	902(42.9)	
Middle	56(16.9)	372(17.7)	
High	86(25.9)	507(24.1)	
College	48(14.5)	320(15.2)	
Smoking			<.001
No	165(49.7)	1341(63.8)	
Yes	167(50.3)	760(36.2)	
Drinking			.064
No	53(16.0)	427(20.3)	
Yes	279(84.0)	1674(79.7)	
Physical activity(exercise)			.560
No	258(77.7)	1602(76.2)	
Yes	74(22.3)	499(23.8)	
Stress Recognition			<.001
No	118(35.5)	535(25.5)	
Yes	214(64.5)	1566(74.5)	
Taking dietary supplements			.023
No	118(35.5)	617(29.4)	
Yes	214(64.5)	1484(70.6)	
Sedentary time			.507
Less than 5hrs.	85(25.6)	577(27.5)	
Over hrs.	247(74.4)	1524(72.5)	
Oral Health			
Chewing difficulty			.947
Severe	128(38.6)	791(37.7)	
Moderate	54(16.3)	351(16.7)	
None	150(45.1)	959(45.6)	
Periodontitis			.003
No	181(54.5)	1325(63.1)	
Yes	151(45.5)	776(36.9)	

Table 3. depicts the major determinants of metabolic syndrome and the impact of oral among Korean 50 and over aged adults. By multi-variable-logistic-regression explained that gender(male) (OR1.86;95%CI:1.27-2.72; p<.05), not taking dietary supplements (OR1.74;95%CI:1.22-2.48; p<.05) factors were the strongest one of metabolic syndrome. And the next determinant factors were presence of stress(OR1.59;95%CI:1.28-1.98; p<.05), lowest education level(OR1.48;95%CI:0.98-2.21; p<.05), oral health factor(periodontitis) (OR1.39; 95%CI: 1.09-1.78; p<.05), were the major determinants of metabolic syndrome respectively. While the related smoking and drinking factors were showed as not associated one with no significance.

**Table3. The determinant factors of metabolic syndrome and the impact of oral health of 50 and older Koreans by multi-variable logistic regression**

variables	OR	95% CI		p-value
Gender				
Male	1.86	1.27	2.72	.001
Female		Reference		
Age				

50-64	1.28	.98	1.65	.061
65 over		Reference		
Socio-economic status				
Residence				
Urban(Dong)		Reference		
Rural(Eup, Myeon)	0.84	.63	1.13	.262
Marital status(spouse)				
No	1.19	.43	3.24	.730
Yes		Reference		
Income level				
Lowest	1.11	.80	1.55	.522
Middle-low	1.01	.70	1.44	.961
Middle-high	1.01	.71	1.42	.968
Highest		Reference		
Education level				
Elementary	1.48	.98	2.21	.050
Middle	1.16	.74	1.78	.519
High	1.26	.85	1.86	.249
College		Reference		
Health-related status				
Smoking				
No		Reference		
Yes	1.15	.79	1.67	.465
Drinking				
No		Reference		
Yes	1.03	0.74	1.45	.825
Physical activity(exercise)				
No	1.20	.89	1.61	.215
Yes		Reference		
Psychological status				
Stress recognition				
No		Reference		
Yes	1.59	1.28	1.98	.001
Taking dietary supplements				
No	1.74	1.22	2.48	.001
Yes		Reference		
Sedentary time				
Less than 5hrs.		Reference		
Over 5hrs.	1.13	0.86	1.48	.360
Oral Health				
Chewing difficulty				
Severe	.924	.657	1.301	.651
Moderate	.910	.692	1.196	.498
None		Reference		
Periodontitis				

No		Reference		
Yes	1.39	1.09	1.78	.007

These findings suggest that men compared to women, not taking of dietary supplements, much stress, lower education level and periodontitis, as the indicator of oral health, were the major determinants of metabolic syndrome with analyzing 2433 participants aged 50 and older participants of the 6th Korea- National- Health and Nutrition- Examination- Survey. The prevalence of metabolic- syndrome was 13.6%. And this prevalence of metabolic syndrome result was low compared to that of another study (Kim, Y. H., et al, 2018) and these results were thought to be due to the total prevalence regardless of gender. Regarding gender (male), which was found to be the strongest factor of metabolic syndrome in this study, was mainly having common with the results of other studies (Kim, Y. H., et al, 2018; Jang, I., & Kim, J. S., 2019) even though they had some difference in age of participants. But it was not consistent with the results of another studies in case of posttraumatic stress of women who had more metabolic risk (LIhua, M., et al, 2020) and having more metabolic risks in female nurses by an observational study(Chico-Barba, G., et al, 2019). With key regard to dietary supplements, as identified as one of the strongest factor of metabolic syndrome through this study, they have been identified to influence on the metabolic syndrome in several studies (Isola, G., et al, 2019; Moukayed, M., & Grant, W. B., 2019; Lee, T. C., et al, 2014; Wu, G., et al, 2007 ). In addition to that, the cardio-metabolic effect of dietary nitrate was reported to be very critical in the point of view of including mitochondrial respiration in health and disease as well as metabolic effects (Lundberg, J. O., et al, 2018). And the results of this present study on the association between natural agents and oral health like periodontal health were supported by current studies (Jaramillo Flores, M. E., 2019; Isola, G., et al, 2019). Concerning stress, which was identified as the determinant factor of this study, that result was supported in some aspects by another studies(Zahran, A. M., et al, 2019; LIhua, M., et al, 2020). Especially, high stress was also known as being associated with taking the risk of periodontitis (Islam, M. M., et al, 2019). 50-64 aged group compared to 65 and older group was more in being risk of metabolic- syndrome of this study and this result was supported by another study (PEDRO, R. E. L., et al, 2019). In the point of view of periodontitis, as a key indicator of oral health, which was identified as the strong influencing factor of metabolic syndrome of this study, was partially in consistent with the result of another study even though there are difference in the aspects of racial sample and study design (Onubogu, U., et al, 2019) and it was supported by another study which was reported on association between oral health and metabolic- syndrome among elderly (PEDRO, R. E. L., et al, 2019) even though the difference in gender. And that, even the awareness and habits of oral care was reported to be contributed to improving of life satisfaction as well as lifestyles of the elderly (Hirano, M., et al, 2019). And poor oral health like periodontitis was shown to be seriously related with diabetes, one of the metabolic components (Rawal, I., et al, 2019) including peripheral vascular disease (Wang, J., et al, 2019) and these results were partially consistent with our results. Moreover, periodontitis was reported to be associated with occupational stress and low coping condition (Islam, M. M., et al, 2019). Chewing, which was related with tooth loss causing of worsening cognition (Zahran, A. M., et al, 2019; Lee, K. H., & Choi, Y. Y., 2019) was already identified as the indicator of good oral health (Kim, Y. T., et al, 2019; Natarajan, P., et al, 2019) even though it was not significant in this study. In view of education level, as the indicator of socio-economic status, it was suggested the major factor of metabolic syndrome through this study, that was supported by another study which was reported on the close relationship between education level, alcohol consumption and physical activity and metabolic syndrome (Kim, Y. H., Kim, H., & Jee, H., 2018). Smoking and alcohol consumption which were known as the important factors of metabolic syndrome even though they were not significant in this study, were not consistent with the results of other study (Kim, Y. H., Kim, H., & Jee, H., 2018) despite of common results in the aspects of education level and they seem to be cause of gender difference by study design. The strength of this study was based on the nationally representative data with stratified survey in the point of reliability and validity. Major consideration of this study was including and dealing with the importance of the oral health variables such as chewing level and periodontitis as the critical role of potential and influencing on general health of middle- older- adults. The limitation of this study is in cross-sectional nature design. To conclude, gender(male), not taking of dietary supplements, stress as the psychological factor, lower education level and periodontitis as the indicator of oral health were the influencing major determinants of metabolic syndrome. So, public health professionals might be needed to manage the education strategies considering age, gender disparities, education level and promoting oral health to prevent metabolic syndrome. Further researches might be necessary to explore the mechanisms for validity between metabolic syndrome and oral health by gender to prepare early public and nursing interventions in this population.

### Conclusion

In conclusion, it was suggested that gender(male) and not taking of dietary supplements were the major determinants of metabolic syndrome. And then, the next determinant factors were stress of the psychological

status, elementary school in lowest education level and periodontitis in oral health were the important determinants of metabolic syndrome respectively. Therefore, public health and nursing professionals might urgently consider managing the public education strategies heightening on age, gender disparities, dietary habits and improving oral health as well as including to prevent metabolic syndrome. Continued researches should be accomplished to find out the links and associations of metabolic syndrome and dietary habits, psychological areas as well as oral health by gender for the future successful ageing.

#### Funding Support – No funding Support

**Conflict of Interest – The authors declare that they have no conflict of interest**

#### References (Harvard Style)

1. Cervino, G., Terranova, A., Briguglio, F., De Stefano, R., Famà, F., D'Amico, C., ... & Scoglio, C., 2019. Diabetes: Oral health related quality of life and oral alterations. *BioMed research international*, 2019. | <https://doi.org/10.1155/2019/5907195>
2. Kim, J. S., Kim, S. Y., Byon, M. J., Lee, J. H., Jeong, S. H., & Kim, J. B., 2019. Association between periodontitis and metabolic syndrome in a Korean nationally representative sample of adults aged 35–79 years. *International journal of environmental research and public health*, 16(16), 2930.
3. Kaur, K. K., 2019. With the Advancement of Knowledge Regarding Correlation of Oral Health and Obesity Role of Dentist Emphasized to Act in Prevention of Further Progression, along with Association with Pregnancy, Fetal Macrosomia, Beta 3 Adren-ergic Receptor Polymorphisms, Energy Drinks. *EC Dental Science*, 18, pp.1927-1938.
4. Hoyas, I., & Leon-Sanz, M., 2019. Nutritional challenges in metabolic syndrome. *Journal of clinical medicine*, 8(9), pp.1301.
5. Lin, W. T., Lee, C. Y., Tsai, S., Huang, H. L., Wu, P. W., Chin, Y. T., ... & Lee, C. H., 2019. Clustering of metabolic risk components and associated lifestyle factors: A nationwide adolescent study in Taiwan. *Nutrients*, 11(3), pp.584.
6. Islam, M. M., Ekuni, D., Yoneda, T., Yokoi, A., & Morita, M., 2019. Influence of Occupational Stress and Coping Style on Periodontitis among Japanese Workers: A Cross-Sectional Study. *International journal of environmental research and public health*, 16(19), pp.3540.
7. Zahran, A. M., Sayed, S. K., Abd El Hafeez, H. A., Khalifa, W. A., Mohamed, N. A., & Hetta, H. F., 2019. Circulating microparticle subpopulation in metabolic syndrome: relation to oxidative stress and coagulation markers. *Diabetes, metabolic syndrome and obesity: targets and therapy*, 12, pp.485.
8. Lihua, M., Tao, Z., Hongbin, M., Hui, W., Caihong, J., & Xiaolian, J., 2020. Metabolic syndrome risk in relation to posttraumatic stress disorder among trauma-exposed civilians in Gansu Province, China. *Medicine*, 99(1).
9. Jaramillo Flores, M. E. 2019., Cocoa flavanols: Natural agents with attenuating effects on metabolic syndrome risk factors. *Nutrients*, 11(4), pp.751.
10. Isola, G., Polizzi, A., Muraglie, S., Leonardi, R., & Lo Giudice, A., 2019. Assessment of vitamin C and antioxidant profiles in saliva and serum in patients with periodontitis and ischemic heart disease. *Nutrients*, 11(12), pp.2956.
11. Van der Veldena, U., 2020. Vitamin C and Its Role in Periodontal Diseases–The Past and the Present: A Narrative Review. *Oral Health Prev Dent*, 18, pp.115-124.
12. Moukayed, M., & Grant, W. B., 2019. Linking the metabolic syndrome and obesity with vitamin D status: risks and opportunities for improving cardiometabolic health and well-being. *Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy*, 12, pp.1437.
13. Lundberg, J. O., Carlström, M., & Weitzberg, E., 2018. Metabolic effects of dietary nitrate in health and disease. *Cell metabolism*, 28(1), pp.9-22.
14. Lee, T. C., Ivester, P., Hester, A. G., Sergeant, S., Case, L. D., Morgan, T., ... & Chilton, F. H., 2014. The impact of polyunsaturated fatty acid-based dietary supplements on disease biomarkers in a metabolic syndrome/diabetes population. *Lipids in health and disease*, 13(1), pp.196.
15. Wu, G., Collins, J. K., Perkins-Veazie, P., Siddiq, M., Dolan, K. D., Kelly, K. A., ... & Meininger, C. J., 2007. Dietary supplementation with watermelon pomace juice enhances arginine availability and ameliorates the metabolic syndrome in Zucker diabetic fatty rats. *The Journal of nutrition*, 137(12), pp.2680-2685.
16. Onubogu, U., Mansfield, W. M., & Ozbek, I. N., 2019. Oral Healthcare Measures to Improve Overall Health in Older Adults. *J Comp Nurs Res Care*, 4, pp.156.
17. Ismail, F., Muneer, M. U., Ahmed, A. R., Hussain, M. W., Farhan, M., Khan, M. U. Q., & Kashif, M., 2019. FACTORS AFFECTING COGNITION IN INDIVIDUALS WITH TOOTH LOSS. *Annals of Dental Specialty*, Vol, 7(4), pp.7.
18. Rawal, I., Ghosh, S., Hameed, S. S., Shivashankar, R., Ajay, V. S., Patel, S. A., ... & Prabhakaran, D.,

2019. Association between poor oral health and diabetes among Indian adult population: potential for integration with NCDs. *BMC oral health*, 19(1), pp.191.
19. PEDRO, R. E. L., BUGONE, É., DOGENSKI, L. C., CARDOSO, M. Z., HÜBNER DA SILVA, A., LINDEN, M. S. S., ... & VALDEZ, E. J., 2019. Relationship between dentition, anthropometric measurements, and metabolic syndrome in the elderly. *Revista de Odontologia da UNESP*, pp.48.
  20. Kim, Y. T., Choi, J. K., Kim, D. H., Jeong, S. N., & Lee, J. H., 2019. Association between health status and tooth loss in Korean adults: longitudinal results from the National Health Insurance Service-Health Examinee Cohort, 2002–2015. *Journal of Periodontal & Implant Science*, 49(3), pp.158-170.
  21. Kim, Y. H., Kim, H., & Jee, H., 2018. Effects of socioeconomic status, health behavior, and physical activity on the prevalence of metabolic syndrome. *Journal of exercise rehabilitation*, 14(2), pp.183.
  22. Jang, I., & Kim, J. S., 2019. Risk of Cardiovascular Disease Related to Metabolic Syndrome in College Students: A Cross-Sectional Secondary Data Analysis. *International journal of environmental research and public health*, 16(19), pp.3708.
  23. Chico-Barba, G., Jiménez-Limas, K., Sánchez-Jiménez, B., Sámano, R., Rodríguez-Ventura, A. L., Castillo-Pérez, R., & Tolentino, M., 2019. Burnout and metabolic syndrome in female nurses: an observational study. *International journal of environmental research and public health*, 16(11), pp.1993.
  24. Hirano, M., Imai, H., Oya, M., Hirano, M., Imai, H., & Oya, M., 2019. Effect of Awareness and Habits of Oral Care on Life Satisfaction. *Oral health & preventive dentistry*, 17(2), pp.131-138.
  25. Wang, J., Geng, X., Sun, J., Zhang, S., Yu, W., Zhang, X., & Liu, H., 2019. The risk of periodontitis for peripheral vascular disease: a systematic review. *Reviews in cardiovascular medicine*, 20(2), pp.81-89.
  26. Lee, K. H., & Choi, Y. Y., 2019. Association between oral health and dementia in the elderly: a population-based study in Korea. *Scientific reports*, 9(1), pp.1-8.
  27. Natarajan, P., Choudhury, M., Seenivasan, M. K., Jeyapalan, K., Natarajan, S., & Vaidhyanathan, A. K., 2019. Body Mass Index and Tooth Loss: An Epidemiological Study in a Sample of Suburban South Indian Population. *Journal of Pharmacy & Bioallied Sciences*, 11(Suppl 2), S402.