

Bernstein Type Inequalities for Polar Derivative of Polynomial

Barchand Chanam¹

¹Department of Mathematics National Institute of Technology Manipur, Imphal, Manipur, India
 barchand_2004@yahoo.co.in

Article History: Received: 11 January 2021; Accepted: 27 February 2021; Published online: 5 April 2021

Abstract- If $p(z)$ is a polynomial of degree n such that $p(z) \neq 0$ in $|z| < k$, $k \leq 1$, then Govil [Proc. Nat. Acad. Sci., Vol. 50, pp. 50-52, 1980.] proved

$$\max_{|z|=1} |p'(z)| \leq \frac{n}{1+k^n} \max_{|z|=1} |p(z)|,$$

provided $|p'(z)|$ and $|q'(z)|$ attain their maxima at the same point on the circle $|z|=1$, where

$$q(z) = z^n \overline{p\left(\frac{1}{\bar{z}}\right)}.$$

Equality in the above inequality holds for $p(z) = z^n + k^n$.

In this paper, we extend the above inequality and an improved version of this into polar derivative of a polynomial.

Keywords – Polynomial, Polar Derivative of a polynomial, Inequalities, Maximum Modulus.

1. Introduction

It was for the first time, Bernstein [10, 11] investigated an upper bound for the maximum modulus of the first derivative of a complex polynomial on the unit circle in terms the maximum modulus of the polynomial on the same circle and proved the following famous result known as Bernstein’s inequality that if $p(z)$ is a polynomial of degree n , then

$$\max_{|z|=1} |p'(z)| \leq n \max_{|z|=1} |p(z)|. \tag{1.1}$$

Inequality (1.1) is best possible and equality occurs for $p(z) = \lambda z^n$, $\lambda \neq 0$, is any complex number.

If we restrict to the class of polynomials having no zero in $|z| < 1$, then inequality (1.1) can be sharpened as

$$\max_{|z|=1} |p'(z)| \leq \frac{n}{2} \max_{|z|=1} |p(z)|. \tag{1.2}$$

The result is sharp and equality holds in (1.2) for $p(z) = \alpha + \beta z^n$, where $|\alpha| = |\beta|$.

Inequality (1.2) was conjectured by Erdős and later proved by Lax [8].

Simple proofs of this theorem were later given by de-Bruijn [5], and Aziz and Mohammad [2].

It was asked by R.P. Boas that if $p(z)$ is a polynomial of degree n not vanishing in $|z| < k$, $k > 0$, then how large can

$$\left\{ \frac{\max_{|z|=1} |p'(z)|}{\max_{|z|=1} |p(z)|} \right\} \text{ be ?} \tag{1.3}$$

A partial answer to this problem was given by Malik [9], who proved

Theorem A. If $p(z)$ is a polynomial of degree n having no zero in the disc $|z| < k$, $k \geq 1$, then

$$\max_{|z|=1} |p'(z)| \leq \frac{n}{1+k} \max_{|z|=1} |p(z)|. \tag{1.4}$$

The result is best possible and equality holds for $p(z) = (z+k)^n$.

For the class of polynomials not vanishing in $|z| < k, k \leq 1$, the precise estimate for maximum of $|p'(z)|$ on $|z| = 1$, in general, does not seem to be easily obtainable.

For quite some time, it was believed that if $p(z) \neq 0$ in $|z| < k, k \leq 1$, then the inequality analogous to (1.4) should be

$$\max_{|z|=1} |p'(z)| \leq \frac{n}{1+k^n} \max_{|z|=1} |p(z)|. \quad (1.5)$$

till E.B. Saff gave the example $p(z) = \left(z - \frac{1}{2}\right) \left(z + \frac{1}{3}\right)$ to counter this belief.

Govil [6] obtained inequality (1.5) with extra condition. More precisely, he proved the following

Theorem B. If $p(z)$ is a polynomial of degree n such that $p(z) \neq 0$ in $|z| < k, k \leq 1$, then

$$\max_{|z|=1} |p'(z)| \leq \frac{n}{1+k^n} \max_{|z|=1} |p(z)|, \quad (1.6)$$

provided $|p'(z)|$ and $|q'(z)|$ attain their maxima at the same point on the circle $|z| = 1$, where

$$q(z) = z^n \overline{p\left(\frac{1}{\bar{z}}\right)}.$$

Equality in (1.6) holds for $p(z) = z^n + k^n$.

Aziz and Rather [3] further improved the bound of (1.6) by involving $\min_{|z|=k} |p(z)|$.

Theorem C. If $p(z)$ is a polynomial of degree n such that $p(z) \neq 0$ in $|z| < k, k \leq 1$, then

$$\max_{|z|=1} |p'(z)| \leq \frac{n}{1+k^n} \left\{ \max_{|z|=1} |p(z)| - \min_{|z|=k} |p(z)| \right\}, \quad (1.7)$$

provided $|p'(z)|$ and $|q'(z)|$ attain their maxima at the same point on the circle $|z| = 1$, where

$$q(z) = z^n \overline{p\left(\frac{1}{\bar{z}}\right)}.$$

As in Theorem B, equality in (1.6) occurs for $p(z) = z^n + k^n$.

Let $p(z)$ be a polynomial of degree n and α be any real or complex number, the polar derivative of $p(z)$, denoted by $D_\alpha p(z)$, is defined as

$$D_\alpha p(z) = np(z) + (\alpha - z)p'(z). \quad (1.8)$$

The polynomial $D_\alpha p(z)$ is of degree at most $n-1$ and it generalizes the ordinary derivative $p'(z)$ of $p(z)$ in the sense that

$$\lim_{\alpha \rightarrow \infty} \frac{D_\alpha p(z)}{\alpha} = p'(z). \quad (1.9)$$

It is of interest to extend ordinary derivative inequalities into polar derivative of a polynomial, for the later version is a generalization of the first. In this direction, Aziz and Shah [4] for the first time extended (1.1) to polar derivative by proving

Theorem D. If $p(z)$ is a polynomial of degree n then for every real or complex number α with $|\alpha| \geq 1$,

$$\max_{|z|=1} |D_\alpha p(z)| \leq n|\alpha| \max_{|z|=1} |p(z)|. \quad (1.10)$$

Inequality (1.10) becomes equality for $p(z) = az^n$, $a \neq 0$.

Further, Aziz [1] extended inequality (1.2) to polar derivative.

Theorem E. If $p(z)$ is a polynomial of degree n having no zero in the disc $|z| < 1$, then for every real or complex number α with $|\alpha| \geq 1$,

$$\max_{|z|=1} |D_\alpha p(z)| \leq \frac{n}{2} (|\alpha| + 1) \max_{|z|=1} |p(z)|. \tag{1.11}$$

The result is best possible and extremal polynomial is $p(z) = z^n + 1$.

For the class of polynomials not vanishing in the disc $|z| < k$, $k \geq 1$, Aziz [1] obtained the extension of Theorem A to polar derivative of $p(z)$.

Theorem F. If $p(z)$ is a polynomial of degree n having no zero in the disc $|z| < k$, $k \geq 1$, then for every real or complex number α with $|\alpha| \geq 1$,

$$\max_{|z|=1} |D_\alpha p(z)| \leq n \left(\frac{k + |\alpha|}{1 + k} \right) \max_{|z|=1} |p(z)|. \tag{1.12}$$

The result is best possible and equality in (1.12) holds for the polynomial $p(z) = (z + k)^n$, with $\alpha \geq 1$.

In this paper, we extend both the Theorems B and C into polar derivative of a polynomial. More precisely, we prove

Theorem 1. If $p(z)$ is a polynomial of degree n such that $p(z) \neq 0$ in $|z| < k$, $k \leq 1$, then for every real or complex number α with $|\alpha| \geq 1$,

$$\max_{|z|=1} |D_\alpha p(z)| \leq n \frac{(k^n + |\alpha|)}{1 + k^n} \max_{|z|=1} |p(z)|, \tag{1.13}$$

provided $|p'(z)|$ and $|q'(z)|$ attain their maxima at the same point on the circle $|z| = 1$, where

$$q(z) = z^n \overline{p\left(\frac{1}{\bar{z}}\right)}.$$

Dividing both sides of (1.13) by $|\alpha|$ and making limit as $|\alpha| \rightarrow \infty$, we obtain inequality (1.6).

Next, we prove the polar derivative form of Theorem C.

Theorem 2. If $p(z)$ is a polynomial of degree n such that $p(z) \neq 0$ in $|z| < k$, $k \leq 1$, then for every real or complex number α with $|\alpha| \geq 1$,

$$\max_{|z|=1} |D_\alpha p(z)| \leq \frac{n}{1 + k^n} \left\{ (k^n + |\alpha|) \max_{|z|=1} |p(z)| - (|\alpha| - 1) \min_{|z|=k} |p(z)| \right\}, \tag{1.14}$$

provided $|p'(z)|$ and $|q'(z)|$ attain their maxima at the same point on the circle $|z| = 1$, where

$$q(z) = z^n \overline{p\left(\frac{1}{\bar{z}}\right)}.$$

Dividing both sides of (1.14) by $|\alpha|$ and making limit as $|\alpha| \rightarrow \infty$, we get inequality (1.7).

II LEMMA

The following lemma is needed for the proofs of the theorems.

Lemma 2.1. If $p(z)$ is a polynomial of degree n , then on $|z| = 1$,

$$|p'(z)| + |q'(z)| \leq n \max_{|z|=1} |p(z)|, \quad (2.1)$$

where

$$q(z) = z^n \overline{p\left(\frac{1}{z}\right)}.$$

The above lemma is a special case of a result due to Govil and Rahman [7].

2. PROOF OF THE THEOREM

I.

Proof of Theorem 1. We omit the proof as it follows on the same lines as that of Theorem 2 by using Theorem B, instead of Theorem C.

Proof of Theorem 2. Let $q(z) = z^n \overline{p\left(\frac{1}{z}\right)}$. Then it is easy to verify that for $|z| = 1$,

$$|q'(z)| = |np(z) - zp'(z)|. \quad (3.1)$$

Now, for every real or complex number α , the polar derivative of $p(z)$ with respect to α is

$$D_\alpha p(z) = np(z) + (\alpha - z)p'(z).$$

This implies for $|z| = 1$,

$$\begin{aligned} |D_\alpha p(z)| &\leq |np(z) - zp'(z)| + |\alpha| |p'(z)| \\ &= |q'(z)| + |\alpha| |p'(z)| \quad (\text{by (3.1)}) \\ &= |q'(z)| + |p'(z)| - |p'(z)| + |\alpha| |p'(z)| \\ &\leq n \max_{|z|=1} |p(z)| + (|\alpha| - 1) |p'(z)| \quad (\text{Lemma 2.1}) \end{aligned} \quad (3.2)$$

Since $|\alpha| \geq 1$, the above inequality when combined with inequality (1.7) of Theorem C gives

$$\begin{aligned} \max_{|z|=1} |D_\alpha p(z)| &\leq n \max_{|z|=1} |p(z)| + (|\alpha| - 1) \left\{ \frac{n}{1+k^n} \left(\max_{|z|=1} |p(z)| - \min_{|z|=k} |p(z)| \right) \right\} \\ &= \frac{n}{1+k^n} \left\{ (k^n + |\alpha|) \max_{|z|=1} |p(z)| - (|\alpha| - 1) \min_{|z|=k} |p(z)| \right\}. \end{aligned}$$

This completes the proof of Theorem 2.

REFERENCES

- A. Aziz, "Inequalities for the polar derivative of a polynomial," J. Approx. theory, vol. 55, pp. b183-193, 1988.
- A. Aziz and Q.G. Mohammad, "Simple proof of a Theorem of Erdős and Lax," Proc. Amer. Math. Soc., vol. 80, pp. 119-122, 1980.
2. Aziz and Rather, "L^f Inequalities for polynomials," Math. Ineq. Appl., vol. 1, pp. 177-191, 1998.
- A. Aziz and W.M. Shah, "Inequalities for the polar derivative of a polynomial," Indian J. Pure and Appl. Math. Vol. 29(2), pp. 163-173, 1998.
4. N.G. de-Bruijn, "Inequalities concerning polynomials in the complex domain, Nederl. Akad. Wetench. Proc. Ser. A, vol. 50(1947), pp. 1265-1272, 1947, Indag. Math., vol. 9, pp. 591-598, 1947.
6. N.K. Govil, "On the Theorem of S. Bernstein," Proc. Nat. Acad. Sci., vol. 50, pp. 50-52, 1980.
7. N.K. Govil and Q.I. Rahman, "Functions of exponential type not vanishing in a half-plane and related polynomials," Trans. Amer. Math. Soc., vol. 137, pp. 501-517, 1969.
8. P.D. Lax, "Proof of a conjecture of P. Erdős on the derivative of a polynomial," Bull. Amer. Math. Soc., vol. 50, pp. 509-513, 1944.

9. M.A. Malik, " On the derivative of a polynomial, " J. London Math. Soc., vol. 1, pp. 57-60, 1969.
10. G.V. Milovanovic, D.S. Mitrinovic and Th. M. Rassias, " Topics in polynomials, Extremal properties, Inequalities, Zeros, " World Scientific Publishing Co., Singapore, 1994.
11. C. Schaeffer, " Inequalities of A. Markoff and S. N. Bernstein for polynomials and related functions, " Bull, Amer. Math. Soc., pp. 565- 579, 1941.